

Data reported as: % susceptible (# isolates tested)¹

	B bron	E coli	E fael	E faem	Ente	K pneu	P aer	P mult	Pseu	S aur	S can	S pint
Amikacin	100% (2)	100%(37)	29% (17)	25% (4)	86% (7)	100% (1)	100%(13)	100% (5)	100% (5)	100% (7)	0% (5)	100% (6)
Amoxicillin/Clavulanic Acid	100% (2)	84% (37)	94% (17)	50% (4)	29% (7)	100% (1)	0% (13)	100% (5)	20% (5)	100% (7)	100% (5)	100% (6)
Ampicillin	50% (2)	68% (37)	94% (17)	50% (4)	57% (7)	0% (1)	8% (13)	100% (5)	20% (5)	43% (7)	100% (5)	67% (6)
Cefazolin	0% (2)	89% (37)	6% (17)	25% (4)	29% (7)	100% (1)	0% (13)	100% (5)	0% (5)	100% (7)	100% (5)	100% (6)
Cefovecin	ND	94% (16)	0% (13)	0% (2)	71% (7)	100% (1)	0% (10)	100% (4)	0% (5)	100% (4)	100% (3)	100% (2)
Cefoxitin	0% (2)	89% (37)	6% (17)	0% (4)	29% (7)	100% (1)	0% (13)	100% (5)	40% (5)	100% (7)	100% (5)	100% (6)
Cefpodoxime	0% (2)	86% (37)	18% (17)	0% (4)	71% (7)	100% (1)	0% (13)	100% (5)	0% (5)	100% (7)	100% (5)	100% (6)
Ceftiofur	0% (2)	86% (37)	6% (17)	0% (4)	71% (7)	100% (1)	0% (13)	100% (5)	20% (5)	100% (7)	100% (5)	100% (6)
Cephalothin	0% (2)	71% (21)	0% (4)	50% (2)	ND	ND	0% (3)	100% (1)	ND	100% (3)	100% (2)	100% (4)
Chloramphenicol	100% (2)	86% (37)	94% (17)	100% (4)	71% (7)	100% (1)	8% (13)	100% (5)	20% (5)	57% (7)	100% (5)	100% (6)
Clindamycin	0% (2)	0% (37)	12% (17)	25% (4)	0% (7)	0% (1)	0% (13)	0% (5)	0% (5)	100% (7)	80% (5)	67% (6)
Doxycycline	ND	94% (16)	92% (13)	100% (2)	71% (7)	100% (1)	20% (10)	100% (4)	40% (5)	100% (4)	67% (3)	0% (2)
Enrofloxacin	50% (2)	78% (37)	35% (17)	0% (4)	86% (7)	100% (1)	69% (13)	100% (5)	100% (5)	71% (7)	40% (5)	50% (6)
Erythromycin	0% (2)	0% (37)	12% (17)	0% (4)	0% (7)	0% (1)	0% (13)	0% (5)	0% (5)	43% (7)	0% (5)	83% (6)
Gentamicin	100% (2)	95% (37)	53% (17)	25% (4)	86% (7)	100% (1)	92% (13)	100% (5)	100% (5)	100% (7)	60% (5)	100% (6)
Imipenem	100% (2)	100%(37)	94% (17)	50% (4)	86% (7)	100% (1)	100%(13)	100% (5)	100% (5)	100% (7)	100% (5)	100% (6)
Marbofloxacin	NI	89% (37)	55% (17)	0% (4)	86% (7)	100% (1)	100%(13)	100% (5)	100% (5)	100% (7)	100% (5)	67% (6)
Orbifloxacin	100% (2)	90% (21)	0% (4)	0% (2)	ND	ND	33% (3)	100% (1)	ND	100% (3)	0% (2)	75% (4)
Oxacillin ³	NA	NA	NA	NA	NA	NA	NA	NA	NA	100% (7)	NA	100% (6)
Penicillin	0% (2)	0% (37)	94% (17)	50% (4)	0% (7)	0% (1)	0% (13)	60% (5)	0% (5)	43% (7)	100% (5)	67% (6)
Rifampin	100% (2)	0% (37)	29% (17)	100% (4)	NI	NI	0% (13)	NI	NI	NI	NI	100% (6)
Tetracycline	100% (2)	86% (21)	50% (4)	50% (2)	ND	ND	0% (3)	100% (1)	ND	100% (3)	50% (2)	75% (4)
Ticarcillin	100% (2)	70% (37)	18% (17)	25% (4)	71% (7)	0% (1)	92% (13)	100% (5)	100% (5)	100% (7)	100% (5)	100% (6)
Ticarcillin/Clavulanic Acid	100% (2)	86% (37)	18% (17)	25% (4)	71% (7)	100% (1)	92% (13)	100% (5)	100% (5)	100% (7)	100% (5)	100% (6)
Trimethoprim/Sulphamethoxazole	50% (2)	92% (37)	88% (17)	100% (4)	86% (7)	100% (1)	15% (13)	100% (5)	80% (5)	100% (7)	100% (5)	67% (6)

³ Isolates resistant to oxacillin are interpreted as methicillin resistant.

Key:

1	Data is reported as: % susceptible (# isolates tested) - not all bacteria isolated at ISU VDL have been tested for antimicrobial susceptibility	
2	See Salmonella serotype table for most common serotypes isolated within each group	
3	Isolates resistant to oxacillin are interpreted as potentially methicillin resistant.	
4	A result of ≤ 2 ug/ml for Carbadox is a conservative indicator of bacterial inhibition by this antimicrobial agent. The result shown is based on pharmacokinetic research indicating an average Carbadox level of 4.5 mcg/ml in the small intestine of pigs fed a dose rate of 50 g/ton. (De Graff 1988).	
5	Multidrug resistant isolates were found resistant to most classes of antimicrobial in the 1 st round of testing. This table represents additional Disk Diffusion testing for those isolates.	
NA	Not applicable	
ND	Not done	
NI	No interpretation	
A equ - Actinobacillus equuli	H ecol - hemolytic E. coli	S aur - Staphylococcus aureus
A suis - Actinobacillus suis	H som - Histophilus somni	S beta- Beta Streptococcus species
Abua - Acinetobacter species	HPS - Haemophilus parasuis	S can - Streptococcus canis
Amy - Actinomyces species	K pneu - Klebsiella pneumoniae	S chol - Salmonella choleraesuis
APP - Actinobacillus pleuropneumoniae	M bov - Moraxella bovis	S dysg - Streptococcus dysgalactiae
B bron - Bordetella bronchiseptica	M haem - Mannheimia haemolytica	S epi- Staphylococcus epidermidis
B tre - Bibersteinia trehalosi (formerly Pasteurella trehalosi)	P aer - Pseudomonas aeruginosa	S equi - Streptococcus equi
Bact - Bacteroides group	P cab - Pasteurella caballi	S equus - Streptococcus equisimilis
C diff - Clostridium difficile	P mult - Pasteurella multocida	S pint - Staph pseudintermedius
C perf - Clostridium perfringens	Past - Pasteurella species	S suis - Streptococcus suis
Clos - Clostridium species	Pec - Peptococcus species	S ube - Streptococcus uberis
E coli - Escherichia coli	Pes - Peptostreptococcus species	S zoo - Streptococcus zooepidemicus
E fael - Enterococcus faecalis	Pmul A - Pasteurella multocida Type A	Salm sp- Salmonella species
E faem - Enterococcus faecium	Pmul D - Pasteurella multocida Type D	Salm B - Salmonella species group B
Enc - Enterococcus species	Prot - Proteus species	Salm C1 - Salmonella species group C1
Ente - Enterobacter species	Prp - Propionibacterium species	Salm C2 - Salmonella species group C2
Erys - Erysipelothrix	Pseu - Pseudomonas species	Salm D - Salmonella species group D
Fus - Fusobacterium	R equ - Rhodococcus equi	Salm E - Salmonella species group E
G ana - Gallibacterium anatis		