

Data reported as: % susceptible (# isolates tested)¹

Antibiotic	B tre	E coli	M haem	P mult	Salm sp
Ampicillin	100% (5)	45% (33)	98% (44)	88% (8)	89% (18)
Ceftiofur	100% (5)	82% (33)	100% (44)	100% (8)	94% (18)
Chlortetracycline	40% (5)	12% (33)	95% (44)	88% (8)	83% (18)
Clindamycin	0% (5)	0% (33)	0% (44)	0% (8)	0% (18)
Danofloxacin	80% (5)	79% (33)	93% (44)	88% (8)	72% (18)
Enrofloxacin	100% (5)	82% (33)	98% (44)	88% (8)	94% (18)
Florfenicol	100% (5)	3% (33)	100% (44)	100% (8)	78% (18)
Gentamicin	100% (5)	79% (33)	100% (44)	100% (8)	94% (18)
Neomycin	100% (5)	45% (33)	89% (44)	88% (8)	100% (18)
Oxytetracycline	40% (5)	12% (33)	82% (44)	75% (8)	83% (18)
Penicillin	0% (5)	0% (33)	39% (44)	75% (8)	0% (18)
Spectinomycin	0% (5)	3% (33)	93% (44)	100% (8)	0% (18)
Sulfadimethoxine	100% (5)	18% (33)	57% (44)	13% (8)	6% (18)
Tiamulin	100% (5)	0% (33)	95% (44)	100% (8)	0% (18)
Tilmicosin	100% (5)	0% (33)	98% (44)	100% (8)	0% (18)
Trimethoprim/Sulphamethoxazole	100% (5)	64% (33)	5% (44)	0% (8)	100% (18)
Tulathromycin	NI	NI	93% (44)	100% (8)	0% (18)